

REFERENCES

1. Armijo-Olivo S, Pitance L, Singh V, Neto F, Thie N, Michelotti A. Effectiveness of Manual Therapy and Therapeutic Exercise for Temporomandibular Disorders: Systematic Review and Meta-Analysis. *Phys Ther* . 2015;96(1):9-25. doi:10.2522/ptj.20140548
2. Fadol Y, Leresche L, Mancl L, Heaton L, Truelove E. Referred Pain in Temporomandibular Disorders: Prevalence, Associated Factors and Effects on TMD Prognosis. *Oral Surg Oral Med Oral Pathol Oral Radiol* . 2018;126(4):e205-e206. doi:10.1016/j.oooo.2018.05.041
3. Friscia M, Sbordone C, Petrocelli M et al. Complications after orthognathic surgery: our experience on 423 cases. *Oral Maxillofac Surg* . 2017;21(2):171-177. doi:10.1007/s10006-017-0614-5
4. List T, Jensen R. Temporomandibular disorders: Old ideas and new concepts. *Cephalalgia* . 2017;37(7):692-704. doi:10.1177/0333102416686302
5. Mercuri L. Temporomandibular Joint Disorder Management in Oral and Maxillofacial Surgery. *Journal of Oral and Maxillofacial Surgery* . 2017;75(5):927-930. doi:10.1016/j.joms.2016.10.033
6. Wieckiewicz M, Boening K, Wiland P, Shiau Y, Paradowska-Stolarz A. Reported concepts for the treatment modalities and pain management of temporomandibular disorders. *Journal of Headache and Pain* . 2015;16(1). doi:10.1186/s10194-015-0586-5
7. Zhang C, Wu JY, Deng DL, et al. Efficacy of Splint Therapy for the Management of Temporomandibular Disorders: a Meta-Analysis. *Oncotarget* . 2016;7(51):84043–84053. doi:10.18632/oncotarget.13059
8. Conti PC, Corrêa AS, Lauris JR, Stuginski-Barbosa J. Management of painful temporomandibular joint clicking with different intraoral devices and counseling: a controlled study. *J Appl Oral Sci*. 2015;23(5):529-35.
9. Costa YM, Porporatti AL, Stuginski-barbosa J, Bonjardim LR, Conti PC. Additional effect of occlusal splints on the improvement of psychological aspects in temporomandibular disorder subjects: A randomized controlled trial. *Archives of Oral Biology*. 2015;60(5):738-44.
10. Gomes CA, Politti F, Andrade DV, et al. Effects of massage therapy and occlusal splint therapy on mandibular range of motion in individuals with temporomandibular

disorder: a randomized clinical trial. *Journal of Manipulative Physiological Therapeutics*. 2014;37(3):164-9. doi: 10.1016/j.jmpt.2013.12.007.

11. Qvintus V, Suominen AL, Huttunen J, Raustia A, Ylöstalo P, Sipilä K. Efficacy of stabilisation splint treatment on facial pain - 1-year follow-up. *Journal of Oral Rehabilitation*. 2015;42(6):439-446. doi:10.1111/joor.12275.
12. Nagata K, Maruyama H, Mizuhashi R, et al. Efficacy of stabilisation splint therapy combined with nonsplint multimodal therapy for treating RDC/TMD axis I patients: a randomised controlled trial. *Journal of Oral Rehabilitation*. 2015;42(12):890-899. doi:10.1111/joor.12332.
13. Kokkola O, Suominen AL, Qvintus V, et al. Efficacy of stabilisation splint treatment on the oral health-related quality of life-A randomised controlled one-year follow-up trial. *Journal of Oral Rehabilitation*. 2018;45(5):355-362.
14. Van Grootel RJ, Buchner R, Wismeijer D, van der Glas HW. Towards an optimal therapy strategy for myogenous TMD, physiotherapy compared with occlusal splint therapy in an RCT with therapy-and-patient-specific treatment durations. *BMC Musculoskeletal Disorders*. 2017;18(1):76. doi:10.1186/s12891-017-1404-9.