

References

1. Cakit BDM, Nacir BM, Genç HM, et al. Cycling progressive resistance training for people with multiple sclerosis: A randomized controlled study. *AM J PHYS MED REHAB.* 2010;89(6):446-457.
2. Castro-Sanches AM, Mataran-Penarrocha G, et al. Hydrotherapy for the treatment of pain in people with multiple sclerosis: A randomized controlled trial. *Evid-Based Compl Alt.* 2012; 2012:8 pages
3. Karagarfard MP, Shariat AP, Ingle LP, Cleland JP, Kargarfard MM. Randomized controlled trial to examine the impact of aquatic exercise training on functional capacity, balance, and perceptions of fatigue in female patients with multiple sclerosis. *Arch Phys Med Rehab.* 2018;99(2):234-241.
4. Kierkegaard M, Lundberg I, Olsson T, et al. High-intensity resistance training in multiple sclerosis — an exploratory study of effects on immune markers in blood and cerebrospinal fluid, and on mood, fatigue, health-related quality of life, muscle strength, walking and cognition. *J Neurol Sci.* 2016;362:251-257.
5. Kooshiar H, Moshtagh M, Sardar M, Foroughipour M, Shakeri M, Vahdatinia B. Fatigue and quality of life of women with multiple sclerosis: A randomized controlled clinical trial. *J Sport Med Phys Fit.* 2015;55(6):668-674
6. Manca A, Dvir Z, Dragone D, Mureddu G, Bua G, Deriu F. Time course of strength adaptations following high-intensity resistance training in individuals with multiple sclerosis. *Eur J Appl Physiol.* 2017;117(4):731-743.

7. Razazian N, Yavari Z, Farnia V, et al. Exercising impacts on fatigue, depression, and paresthesia in female patients with multiple sclerosis. *Med Sci Sport Exer.* 2016;48(5):796-803.